

ARTWORK DESIGN STANDARD INSERT

Date of use: 16/10/2025
(DD-MM-YYYY)

MERSIDOL
Size (L x W) 159 x 99 mm

Material
HVS Paper 60g
Packsize/Primary
Syrup 120 mg/5 mL
Primary
Plastic Bottle @ 62 mL
No. Reg
MERSIDOL ANAK, No. Reg. XXXXXXXXXXXX

Front / Back	
Type Font	
Product Name/ Active ingredient	Another Font
Insaniburger Regular	Arial Regular / Bold
9.89 & 7,9 pt	6pt

	
16 x 4mm (LxH)	TEXT Black
Material Code	
IIN00351	

MERSIDOL®
PARACETAMOL
Sirup 120 mg/5 mL

PEMERIAN OBAT
Larutan jernih, warna merah, rasa manis sedikit pahit aroma strawberry.

KOMPOSISI
Tiap 5 mL sirup mengandung:
Paracetamol 120 mg

ZAT TAMBAHAN
Propylene Glycol, Saccharin sodium, Povidone K-30, Strawberry flavour, Sodium benzoate, Sodium citrate dihydrate, Purified water, Citric acid anhydrate, Carmoisine CI 14720.

CARA KERJA
Analgesik-antipiretik
- Sebagai analgesik, bekerja dengan meningkatkan ambang rangsang rasa sakit.
- Sebagai antipiretik, diduga bekerja langsung pada pusat pengatur panas di hipotalamus.

INDIKASI
Meringankan rasa sakit pada keadaan sakit kepala, sakit gigi dan menurunkan demam.

POSOLOGI DAN CARA PEMBERIAN
Posologi
1-2 tahun : 1 sendok takar (5 mL) 3-4 kali sehari
2-6 tahun : 1-2 sendok takar (5-10 mL) 3-4 kali sehari
6-9 tahun : 2-3 sendok takar (10-15 mL) 3-4 kali sehari
9-12 tahun : 3-4 sendok takar (15-20 mL) 3-4 kali sehari
atau sesuai petunjuk dokter.
Cara Pemberian
Obat diberikan secara oral.

KONTRAINDIKASI
- Penderita gangguan fungsi hati yang berat.
- Penderita hipersensitif terhadap obat ini.

EFEK SAMPING
- Penggunaan jangka lama dan dosis besar dapat menyebabkan kerusakan hati.
- Reaksi hipersensitivitas.

PERINGATAN DAN PERHATIAN
- Hati-hati penggunaan obat ini pada penderita penyakit ginjal.
- Bila setelah 2 hari demam tidak menurun atau setelah 5 hari nyeri tidak menghilang segera hubungi Unit Pelayanan Kesehatan.
- Penggunaan obat ini pada penderita yang mengkonsumsi alkohol, dapat meningkatkan risiko kerusakan fungsi hati.

CARA PENYIMPANAN
Simpan di bawah suhu 30°C, terlindung dari cahaya.
Jauhkan obat dari jangkauan anak-anak.
Perhatian : Setelah kemasan dibuka, Mersidol Sirup tidak boleh disimpan lebih dari 30 hari.

KEMASAN DAN NOMOR REGISTRASI
Dus, botol plastik @ 60 mL MERSIDOL sirup, No. Reg. XXXXXXXXXXXXXXXX

Untuk pelaporan Efek Samping dan Keluhan Produk, dapat menghubungi:
Tim Farmakovigilans PT. Mersifarma
Email : farmakovigilans@mersifarma.com
Telepon : 0818-8287-0825

KOCOK DAHULU SEBELUM DIGUNAKAN

Diproduksi oleh:

PT. Mersifarma TM
Sukabumi – Indonesia

MERSIDOL®
PARACETAMOL
Syrup 120 mg/5 mL

DRUG DESCRIPTION
Clear solution, red color, sweet taste slightly bitter strawberry flavour.

COMPOSITION
Each 5 mL syrup contains:
Paracetamol 120 mg

EXCIPIENTS
Propylene Glycol, Saccharin sodium, Povidone K-30, Strawberry flavour, Sodium benzoate, Sodium citrate dihydrate, Purified water, Citric acid anhydrate, Carmoisine CI 14720.

ACTIONS
Analgesic-antipyretic
- As an analgesic, works by increasing the pain threshold.
- As an antipyretic, by direct action on the hypothalamic heat regulating center.

INDICATIONS
To relieve pain in headaches, toothaches, and reduce fever.

POSODOLOGY AND METHOD OF ADMINISTRATION
Posology
1-2 years : 1 spoon (5 mL), 3-4 times a day
2-6 years : 1-2 spoon (5-10 mL), 3-4 times a day
6-9 years : 2-3 spoon (10-15 mL), 3-4 times a day
9-12 years : 3-4 spoon (15-20 mL), 3-4 times a day
Or as directed by the physician.
Method of Administration
The drug is given orally.

CONTRAINDICATIONS
- Patients with severe hepatic dysfunctions.
- Patients with hypersensitivity to this medicine.

SIDE EFFECTS
- Prolonged use and overdosage may cause liver damage.
- Hypersensitivity reactions.

WARNINGS AND PRECAUTIONS
- Should be used with caution in patients with renal disease.
- Consult to Health Service Unit if fever persists for more than 2 days or if pain continuous for more than 5 days.
- The use of this drug in patient consuming alcohol may increase the risk of liver damage.

STORAGE
Store at temperature below 30°C, protect from light.
Keep out of the reach of children.
Precaution: After opened, Mersidol syrup should not be stored for more than 30 days.

PRESENTATION AND REGISTRATION NUMBER
Box, plastic bottle @ 60 mL Syrup of MERSIDOL, Reg. No XXXXXXXXXXXXXXXX

For reporting Side Effects and Product Complaints, please contact:
PT. Mersifarma Pharmacovigilance Team
Email : farmakovigilans@mersifarma.com
Telephone : 0818-8287-0825

SHAKE WELL BEFORE USE

Manufactured by:

PT. Mersifarma TM
Sukabumi – Indonesia
000000-000000-00

159

IIN00351

99